

# SAFESTORE: Instructions

## SAWTOOTH/MERCHANT GRAIN BEETLE

### *Oryzaephilus spp.*

#### *With Easy Read Traps*

*Oryzaephilus* spp. monitoring kit contains ten “Easy Read” units, 10 rubber septas containing kairomone (food attractant), and instructions. A chart for record keeping may be obtained from our website: [www.jfoakes.com](http://www.jfoakes.com)

Best results can be obtained by using the SAFESTORE system to set up a monitoring program. When in place, such a program can help you to identify when and where infestation problems will arise.

**RECOMENDED:** that a thorough inspection of the area involved be carried out, and potential infestation “hotspots” are identified and marked on a site plan or map. The position of the traps can be marked on this map when they are placed, to facilitate the reading of catch levels.

**PREPARATION:** Simply place 1 rubber septa into the center of the glue area inside the Easy Read trap. It is now ready to place.

**PLACEMENT:** can affect the amount of insects that will be caught, so for an effective program, it is important that the traps are placed in the best position available, and when they are replaced, the positioning is altered as little as possible so that information from different times of the year can be compared. Traps should be placed when temperatures reach 55<sup>o</sup>F or higher.

**BEST POSITIONING:** varies from site to site, so there is a certain amount of choice available in the placing of the traps, however good results can be obtained by following a few guidelines:

- Wherever possible, place units on a flat even surface.
- Sheltered and recessed areas provide good monitoring points, ex: fire hoses or fire extinguishers.
- Ensure that sanitation staff is informed of the program to prevent trap removal.
- Mark the position of the traps on the site plan, and assign them a number.
- Never store monitoring equipment with insecticides.
- Wash hands before placing or inspecting trap units.
- Avoid placing traps in areas where large volumes of air are moving out of the building.
- Place traps in a grid pattern (30 – 50 feet) or shorter intervals to pinpoint infestations.

**REGULAR CHECKING:** once per week is recommended, however it may be necessary to inspect more often if you have a zero insect tolerance policy.

The sensitivity of the area to be monitored dictates how often they should be inspected, but whatever frequency they are checked should be kept constant so that the records you keep can be compared to each other. [Click here](#) to see our Record Charts for monitoring of specific and non-target pests. These tables can be printed, photocopied, completed and filed for future reference.

Trap units should be replaced every 4 - 6 weeks. Care should be taken during inspections to check the condition of the glue areas in the units, especially in dusty conditions or high insect catch situations, which may cause the glue surface to deteriorate. Should this occur, the trap should be replaced.

Un-used lures/pheromones and traps should be stored in a cool area, avoid direct sunlight. Lures/pheromones may be refrigerated for longer shelf life.

The information given in this instruction sheet is provided as a general guide, and is by no means extensive. The biology of pests is the subject of a great many texts and although every effort has been made to provide factually correct information, Russell Fine Chemicals nor J.F.Oakes Sales & Marketing will in no circumstance be liable in respect of any omission or error.